

Ricci Flow Equations on Special Finsler Space

S. K. Narasimhamurthy and D. M. Vasantha

Department of Mathematics, Kuvempu University, Shankaraghatta - 577451,
Shivamogga, Karnataka, INDIA

e-mail: nmurthysk@gmail.com, dmvasantha@gmail.com

(Received: November 11, 2013)

Abstract

In this paper, we deal with one of the special Finsler spaces such as C_2 -like space and find out Ricci flow equations on C_2 -like space with (α, β) -metric.

Key Words: Ricci flow, (α, β) -metric, C_2 -like space.

2010 AMS Subject Classification: 53B40, 53C60.

1. Introduction

Ricci flow is a means by which one can take an arbitrary Riemannian manifold and smooth out the geometry of that manifold to make it look more symmetric. It has proven to be a very useful tool in understanding the topology of such manifolds.

The Ricci flow theory became a very powerful method in understanding the geometry and topology of Riemannian manifolds ([3], [7]-[9]). The most important achievement of this theory was the geometrization conjecture of Thurston. One consequence of this conjecture is the Poincare conjecture. This conjecture was formulated by Henri Poincare [4] and proved by Perelman ([7]-[9]). The proof of Poincare conjecture based on a detailed analysis of Ricci flow surgery is one of the most impressive recent achievement of modern mathematics.

S. Vacaru ([12]-[18]) studied on nonholonomic Ricci flows, evolution equations and dynamics, exact solutions in gravity, symmetric and non symmetric metrics, the entropy of Lagrange-Finsler spaces and Ricci flows, spectral functionals, nonholonomic Dirac operators and non commutative Ricci flows, Fractional nonholonomic Ricci flows, Nonholonomic Ricci flows and parametric deformations of the solitonic pp-waves and schwarzschild solutions. A. Thayebi, E. Peyghan and B. Najafi [11] studied on Ricci flow equation on (α, β) -metrics.

R. S. Hamilton [3] introduced the following geometric evolution equation for a Riemannian metric g_{ij} and the corresponding Ricci curvature tensor Ric_{ij}

$$\frac{d}{dt}(g_{ij}) = -2Ric_{ij}, \quad g(t=0) = g_0 \quad (1.1)$$

is known as the un-normalized Ricci flow in Riemannian geometry. Hamilton showed that there is a unique solution to this equation for an arbitrary smooth metric on a closed manifold over a sufficiently short time.

In this paper, we deal with one of the special Finsler spaces such as C_2 -like space and find out un-normal Ricci flow and normal Ricci flow equations on C_2 -like space with (α, β) -metric.

2. Preliminaries

Definition 2.1. A Finsler metric is a scalar field $L(x, y)$ which satisfies the following three conditions:

- (i) It is defined and differential for any point of $TM^n \setminus \{0\}$,
- (ii) It is positively homogeneous of first degree in y^i , that is,

$$L(x, \lambda y) = \lambda L(x, y), \quad \text{for any positive number } \lambda,$$
- (iii) It is regular, that is,

$$g_{ij}(x, y) = \frac{1}{2} \dot{\partial}_i \dot{\partial}_j L^2,$$
 constitute the regular matrix g_{ij} , where $\dot{\partial}_i = \frac{\partial}{\partial y^i}$.

The manifold M^n equipped with a fundamental function $L(x, y)$ is called Finsler space $F^n = (M^n, L)$.

The concept of the (α, β) -metric was introduced in 1972 by M. Matsumoto and has been studied by M. Hashiguchi, Y. Ichijyo, S. Kikuchi, C. Shibata and others ([5], [6] and [10]).

Definition 2.2. The Finsler space $F^n = (M^n, L)$ is said to have an (α, β) -metric if L is a positively homogeneous function of degree one in two variables $\alpha = \sqrt{a_{ij}y^i y^j}$ and $\beta = b_i(x)y^i$, where α is a Riemannian metric and β is differential 1-form.

A deformation of Finsler metrics means a 1-parameter family of metrics $g_{ij}(x, y, t)$, such that $t \in [-\epsilon, \epsilon]$ and $\epsilon > 0$ is sufficiently small. For such a metric $w = u_i dx^i$, the volume element as well as the connections attached to it depend on t . The same equation can be used in the Finsler setting. Another Ricci flow equation can also be used instead of this tensor evolution equation [2]. By contracting $\frac{d}{dt}g_{ij} = -2Ric_{ij}$ with y^i and y^j gives, via Euler's theorem,

we get

$$\frac{\partial L^2}{\partial t} = -2L^2R,$$

where $R = \frac{1}{L^2} Ric$. That is,

$$d \log L = -R, \quad L(t = 0) = L_0.$$

This scalar equation directly addresses the evolution of the Finsler metric L and makes geometrical sense on both the manifold of nonzero tangent vectors TM_0 and the manifold of rays. It is therefore suitable as an un-normalized Ricci flow for Finsler geometry.

By using the elegance work of Akbar-Zadeh in [1], Bao proposed the following normalised Ricci flow equation for Finsler metrics

$$\frac{d}{dt} \log L = -R + \frac{1}{Vol(SM)} \int_{SM} R dV, \quad L(t = 0) = L_0,$$

where the underlying manifold M is compact [2].

It is noted that [11], Chern had asked whether every smooth manifold admits a Ricci-constant Finsler metric? The weaker case of this question is that whether every smooth manifold admits a Einstein Finsler metric? His question has already been settled in the affirmative for dimension 2 because, by a construction of Thurstons, every Riemannian metric on a two-dimensional manifold admits a complete Riemannian metric of constant Gaussian curvature.

Let M be an n -dimensional C^∞ manifold, T_xM be the tangent space at $x \in M$ and $TM = \cup_{x \in M} T_xM$ be the tangent bundle of M . Let $x \in M$ and $L_x = L|_{T_xM}$. To measure the non-Euclidean feature of L_x , define $C_y : T_xM \otimes T_xM \otimes T_xM \rightarrow \mathbb{R}$ by

$$C_y(u, v, w) = \frac{1}{2} \frac{d}{dt} [g_{y+tw}(u, v)]|_{t=0}, \quad u, v, w \in T_xM.$$

The family $C = \{C_y\}_{y \in TM_0}$ is called the Cartan torsion. It is known that $C = 0$ if and only if L is Riemannian.

For $y \in T_xM_0$, define mean Cartan torsion I_y by $I_y(u) = I_i(y)u^i$, where $I_i = g^{jk}C_{ijk}$, $C_{ijk} = \frac{1}{2} \frac{\partial g_{ij}}{\partial y^k}$ and $u = u^i \frac{\partial}{\partial x^i}|_x$. By Deicke's theorem, L is Riemannian if and only if $I_y = 0$.

A Finsler metric L is called C_2 -like if its Cartan tensor is given by

$$C_{ijk} = \frac{1}{\|I\|^2} I_i I_j I_k.$$

3. Un-normal Ricci flow equation on C_2 -like space with (α, β) -metrics

Here, we study (α, β) -metrics satisfying un-normal Ricci flow equation. First, we prove the following lemmas.

Lemma 3.1. Let L_t be a deformation of an (α, β) -metric L , which is C_2 -like, on a manifold M of dimension $n \geq 3$. Then the variation of Cartan tensor is given by the following

$$C'_{ijk} I^i I^j I^k = -2R \|I\|^4 - \frac{1}{2} L^2 R_{,i,j,k} I^i I^j I^k - 3 \|I\|^2 I^m R_{,m}, \quad (3.1)$$

where $\|I\|^2 = I_m I^m$.

Proof. First, assume that L_t be a deformation of a Finsler metric on a two-dimensional manifold M satisfies Ricci flow equation, that is,

$$\frac{d}{dt} g_{ij} = g'_{ij} = -2Ric_{ij}, \quad d \log L = \frac{L'}{L} = -R, \quad (3.2)$$

where $R = \frac{1}{L^2} Ric$. By definition of Ricci tensor, we have

$$\begin{aligned} Ric_{ij} &= \frac{1}{2} [RL^2]_{y^i y^j} \\ &= Rg_{ij} + \frac{1}{2} L^2 R_{,i,j} + R_{,i} y_j + R_{,i} y_i, \end{aligned} \quad (3.3)$$

where $R_{,i} = \frac{\partial R}{\partial y^i}$ and $R_{,i,j} = \frac{\partial^2 R}{\partial y^i \partial y^j}$. Taking a vertical derivative of (3.3) and using $y_{i,j} = g_{ij}$ and $LL_k = y_k$ yields

$$\begin{aligned} Ric_{ij,k} &= 2RC_{ijk} + \frac{1}{2} L^2 R_{,i,j,k} + \{g_{jk} R_{,i} + g_{ij} R_{,k} + g_{ki} R_{,j}\} \\ &\quad + \{R_{,j,k} y_i + R_{,i,j} y_k + R_{,k,i} y_j\}. \end{aligned} \quad (3.4)$$

Contracting (3.4) with $I^i I^j I^k$ and using $y_i I^i = y^i I_i = 0$ implies that

$$Ric_{ij,k} I^i I^j I^k = 2RC_{ijk} I^i I^j I^k + \frac{1}{2} L^2 R_{,i,j,k} I^i I^j I^k + 3 \|I\|^2 I^m R_{,m}. \quad (3.5)$$

The Cartan tensor of an (α, β) -metric on n -dimensional manifold M is given by

$$C_{ijk} = \frac{1}{\|I\|^2} I_i I_j I_k. \quad (3.6)$$

Multiplying (3.6) with $I^i I^j I^k$ yields

$$C_{ijk} I^i I^j I^k = \|I\|^4. \quad (3.7)$$

Then by (3.5) and (3.7), we get

$$\begin{aligned} Ric_{ij,k} I^i I^j I^k &= 2R \|I\|^4 + \frac{1}{2} L^2 R_{,i,j,k} I^i I^j I^k \\ &\quad + 3 \|I\|^2 I^m R_{,m}. \end{aligned} \quad (3.8)$$

On the other hand, since L_t satisfies Ricci flow equation, then

$$\begin{aligned} C'_{ijk} &= \frac{1}{2} \frac{\partial g'_{ij}}{\partial y^k} \\ &= \frac{1}{2} \frac{\partial(-2Ric_{ij})}{\partial y^k} \\ &= -Ric_{ij,k}. \end{aligned} \quad (3.9)$$

By (3.8) and (3.9), we get (3.1).

Lemma 3.2. Let L_t be a deformation of an (α, β) -metric L , which is C_2 -like, on a manifold M of dimension $n \geq 3$. Then $C'_{ijk} I^i I^j I^k$ is a factor of $\|I\|^2$.

Proof. Since $g^{ij} g_{jk} = \delta_k^i$, we have

$$\begin{aligned} (g^{ij} g_{jk})' &= 0 \\ \Rightarrow g'^{ij} g_{jk} + g^{ij} g'_{jk} &= 0 \\ \Rightarrow g'^{ij} g_{jk} + g^{ij} (-2Ric_{jk}) &= 0 \\ \Rightarrow g'^{ij} g_{jk} - 2g^{ij} Ric_{jk} &= 0, \end{aligned} \quad (3.10)$$

or equivalently, $(g^{ij})' g_{jk} = 2g^{ij} Ric_{jk}$.

Contracting with g^{lk} gives

$$(g^{il})' = 2Ric^{il}. \quad (3.11)$$

Then, we have

$$\begin{aligned} I'_i &= (g^{jk} c_{ijk})' \\ &= (g^{jk})' c_{ijk} + g^{jk} (c_{ijk})' \\ &= 2Ric^{jk} c_{ijk} + g^{jk} (-Ric_{ij,k}) \\ &= Ric^{jk} \frac{\partial g_{ij}}{\partial y^k} - (g^{jk} Ric_{jk})_{,i} + g^{jk} {}_{,i} Ric_{jk}. \end{aligned} \quad (3.12)$$

Since

$$-g^{jk} Ric_{ij,k} = - (g^{jk} Ric_{jk})_{,i} + g^{jk} {}_{,i} Ric_{jk},$$

we have

$$\begin{aligned}
I'_i &= Ric^{jk} g_{jk,i} - \left(g^{jk} Ric_{jk} \right)_{,i} + g^{jk}_{,i} Ric_{jk} \\
&= - \left(g^{jk} Ric_{jk} \right)_{,i} \\
&= -\rho_i,
\end{aligned} \tag{3.13}$$

where $\rho = g^{jk} Ric_{jk}$ and $\rho_i = \frac{\partial \rho}{\partial y^i}$. Thus

$$\begin{aligned}
I'^i &= (g^{ij} I_j)' \\
&= (g^{ij})' I_j + g^{ij} I'_j \\
&= 2Ric^{ij} I_j + g^{ij} (-\rho_j) \\
&= 2Ric^{ij} I_j - \rho^i.
\end{aligned} \tag{3.14}$$

The variation of $y_i = LL_{y^i}$ with respect to t is given by

$$y'_i = -2Ric_{im} y^m.$$

Therefore, we can compute the variation of angular metric h_{ij} as follows

$$\begin{aligned}
h'_{ij} &= (g_{ij} - L^{-2} y_i y_j)' \\
&= (g_{ij})' - (L^{-2} y_i y_j)' \\
&= -2Ric_{ij} - \left\{ L^{-2} [y'_i y_j + y_i y'_j] + y_i y_j (L^{-2})' \right\} \\
&= -2Ric_{ij} - L^{-2} [-2Ric_{im} y^m y_j - 2Ric_{jm} y^m y_i] - 2L^{-2} R y_i y_j \\
&= -2Ric_{ij} + 2(h_{ij} - g_{ij}) R + 2(Ric_{im} L^{-1} y_j) L^{-1} y^m \\
&\quad + 2(Ric_{jm} L^{-1} y_i) L^{-1} y^m \\
&= -2Ric_{ij} + 2(h_{ij} - g_{ij}) R + 2(Ric_{im} l_j + Ric_{jm} l_i) l^m,
\end{aligned} \tag{3.15}$$

where $l_i = L^{-1} y_i$ and $l^m = L^{-1} y^m$. Thus, we consider the variation of Cartan tensor

$$\begin{aligned}
C'_{ijk} &= \left[\frac{1}{\|I\|^2} I_i I_j I_k \right]' \\
&= \frac{\|I\|^2 (I_i I_j I_k)' - I_i I_j I_k (\|I\|^2)'}{\|I\|^4} \\
&= \frac{(I'_i I_j I_k + I'_j I_i I_k + I'_k I_i I_j)}{\|I\|^2} - \frac{C_{ijk} (I^m I_m + I^m I'_m)}{\|I\|^2} \\
&= -\frac{(\rho_i I_j I_k + \rho_j I_i I_k + \rho_k I_i I_j)}{\|I\|^2} - \frac{(I^m I_m + I^m I'_m) C_{ijk}}{\|I\|^2}.
\end{aligned} \tag{3.16}$$

Multiplying (3.16) with $I^i I^j I^k$ gives

$$\begin{aligned}
C'_{ijk} I^i I^j I^k &= -\frac{(\rho_i I^i + \rho_j I^j + \rho_k I^k) \|I\|^4}{\|I\|^2} \\
&\quad -\frac{(I^m I_m + I^m I'_m)}{\|I\|^2} \times \frac{1}{\|I\|^2} \times I_i I_j I_k \times I^i I^j I^k \\
&= -\frac{3\|I\|^4 \rho_m I^m}{\|I\|^2} - \frac{[(2Ric^{mj} I_j - \rho^m) I_m + I^m (-\rho_m)]}{\|I\|^2} \times \frac{\|I\|^6}{\|I\|^2} \\
&= \|I\|^2 \{ \rho^m I_m - 2(Ric^{mj} I_m I_j + \rho_m I^m) \}, \tag{3.17}
\end{aligned}$$

which implies $C'_{ijk} I^i I^j I^k$ is a factor of $\|I\|^2$. This completes the proof.

Next, we prove the following main theorem.

Theorem 3.1. Suppose that L is an (α, β) -metric on M , which is C_2 -like, then every deformation L_t of the metric L satisfying un-normal Ricci flow equation is an Einstein metric.

Proof. By virtue of lemma and lemma , $R_{,i,j,k} I^i I^j I^k$ is a factor of $\|I\|^2$. Since $R_{,i,j,k} I^i I^j I^k$ is a factor of $\|I\|^2$, multiplying it with y^k or y^j implies $R_{,i} = 0$. It means that $R = R(x)$ and then L_t is an Einstein metric.

4. Normal Ricci flow equation on C_2 -like space with (α, β) -metrics

If M is a compact manifold, then $S(M)$ is compact and we can normalize the Ricci flow equation by requiring that the flow keeps the volume of SM constant. Recalling the Hilbert form $w = L_{y^i} dx^i$, that volume is

$$Vol_{SM} = \int_{SM} \frac{(-1)^{\frac{n(n-1)}{2}}}{(n-1)!} w \wedge (dw)^{n-1} = \int_{SM} dV_{SM}.$$

During the evolution, L , w and consequently the volume form dV_{SM} and the volume Vol_{SM} , all depend on t . On the other hand, the domain of integration SM , being the quotient space of TM_0 under the equivalence relation $z \sim y$, $z = \lambda y$ for some $\lambda > 0$, is totally independent of any Finsler metric and hence does not depend on t . We have

$$\frac{d}{dt}(dV_{SM}) = \left[g_{ij} \frac{d}{dt} g_{ij} - n \frac{d}{dt} \log L \right] dV_{SM}.$$

A normalized Ricci flow for Finsler metrics is proposed by Bao as follows

$$\frac{d}{dt} \log L = -R + \frac{1}{Vol(SM)} \int_{SM} R dV, \quad L(t=0) = L_0, \tag{4.1}$$

where the underlying manifold M is compact. Now, we let $Vol(SM) = 1$. Then all of Ricci constant metrics are exactly the fixed points of the above flow. Let

$$Ric_{ij} = \frac{1}{2} (L^2 R)_{.y^i .y^j}$$

and differentiating (4.1) with respect to y^i and y^j , the following normal Ricci flow tensor evaluation equation is concluded.

$$\frac{d}{dt} g_{ij} = -2Ric_{ij} + \frac{2}{Vol(SM)} \int_{SM} R dV g_{ij}, \quad g(t=0) = g_0. \quad (4.2)$$

Starting with any familiar metric on M as the initial data L_0 , we may deform it using the proposed normalized Ricci flow, in the hope of arriving at a Ricci constant metric.

Theorem 4.1. Suppose that L is an (α, β) -metric on M , which is C_2 -like, then every deformation L_t of the metric L satisfying normal Ricci flow equation is an Einstein metric.

Proof. Consider Finsler surfaces which satisfy the normal Ricci flow equation. Then

$$\begin{aligned} \frac{dg_{ij}}{dt} &= -2Ric_{ij} + 2 \int_{SM} R dV g_{ij}, \\ d \log L &= \frac{L'}{L} = -R + \int_{SM} R dV. \end{aligned} \quad (4.3)$$

By the same argument in the un-normal Ricci flow case, we can calculate the variation of mean Cartan tensor as follows

$$\begin{aligned} I'_i &= (g^{jk} C_{ijk})' \\ &= (g^{jk})' C_{ijk} + g^{jk} (C_{ijk})' \\ &= \left[2Ric^{jk} - 2 \int_{SM} R dV g_{jk} \right] C_{ijk} + g^{jk} \left[Ric_{jk,i} + 2 \int_{SM} R dV C_{ijk} \right] \\ &= -\rho_i. \end{aligned} \quad (4.4)$$

Then we have

$$\begin{aligned} I'^i &= (g^{ij} I_j)' + g^{ij} I'_j \\ &= \left[2Ric^{ij} - 2 \int_{SM} R dV g_{ij} \right] I_j - g^{ij} \rho_j. \end{aligned} \quad (4.5)$$

As the similar way that we used in un-normal Ricci flow, it follows that

$$\begin{aligned} C'_{ijk} &= \left[\frac{1}{\|I\|^2} I_i I_j I_k \right]' \\ &= - \frac{(I^m I_m + I^m I'_m) C_{ijk}}{\|I\|^2} - \frac{(\rho_i I_j I_k + \rho_j I_i I_k + \rho_k I_i I_j)}{\|I\|^2}. \end{aligned} \quad (4.6)$$

Contracting it with $I^i I^j I^k$, we can say $C'_{ijk} I^i I^j I^k$ is a factor of $\|I\|^2$. By lemma , we deduce that $R_{,i,j,k} I^i I^j I^k$ is a factor of $\|I\|^2$. By the same argument, it results that every deformation L_t of the metric L satisfying normal Ricci flow equation is an Einstein metric.

REFERENCES

- [1] **Akbar-Zadeh, H.** : Generalized Einstein manifolds, *J. Geom. Phys.*, 17 (1995), 342-380.
- [2] **Bao, D.** : On two curvature-driven problems in Riemann-Finsler geometry, *Advanced Studies in Pure Mathematics*, xx, (2007).
- [3] **Hamilton, R. S.** : Three manifolds with positive Ricci curvature, *J. Diff. Geom.*, 17 (1982), 255-306.
- [4] **Poincare, Henri** : Cinquieme complement ‘a lanalysis situs. In *OEuvres*. Tome VI, Les Grands Classiques Gauthier-Villars. [Gauthier-Villars Great Classics] pages v+541, Editions Jacques Gabay, Sceaux, Reprint of the 1953 edition, (1996).
- [5] **Narasimhamurthy, S. K., Aveesh, S. T., Nagaraj, H. G. and Pradeep Kumar** : On special hypersurface of a Finsler space with the metric $(\alpha + \frac{\beta^{n-1}}{\alpha^n})$, *Acta Universitatis Apulensis*, 17 (2009).
- [6] **Narasimhamurthy, S. K. and Vasantha, D. M.** : Projective change between two Finsler spaces with (α, β) -metric, *KYUNGPOOK Math. J.*, 52 (1) (2012), 81–89.
- [7] **Perelman, G.** : The entropy formula for the Ricci flow and its geometric applications, arXiv: math.DG/ 0211159.
- [8] **Perelman, G.** : Ricci flow with surgery on three manifolds, arXiv: math.DG/ 03109.
- [9] **Perelman, G.** : Finite extinction time for the solutions to the Ricci flow on certain three manifolds, arXiv: math.DG/ 0307245.
- [10] **Prasad, B. N., Gupta, B. N. and Singh, D. D.** : Conformal transformation in Finsler spaces with (α, β) -metric, *Indian J. Pure and Appl. Math.*, 18 (4) (1961), 290–301.
- [11] **Thayebi, A., Peyghan, E. and Najafi, B.** : Ricci flow equation on (α, β) -metrics, arXiv:1108.0134v1.
- [12] **Vacaru, S.** : Nonholonomic Ricci flows: II. evolution equations and dynamics, *J. Math. Phys.*, 49 (2008), 043504.
- [13] **Vacaru, S.** : Nonholonomic Ricci flows, exact solutions in gravity, and Symmetric and Nonsymmetric Metrics, *Int. J. Theor. Phys.*, 48 (2009), 579-606.
- [14] **Vacaru, S.** : The Entropy of Lagrange-Finsler spaces and Ricci flows, *Rep. Math. Phys.*, 63 (2009), 95-110.

- [15] **Vacaru, S. :** Spectral functionals, nonholonomic Dirac operators, and non commutative Ricci flows, *J. Math. Phys.*, 50 (2009), 073503.
- [16] **Vacaru, S. :** Fractional Nonholonomic Ricci Flows, arXiv:1004.0625.
- [17] **Vacaru, S. :** Nonholonomic Ricci flows: Exact solutions and gravity, *Electronic Journal of Theoretical Physics*, 6, N20 (2009), 27-58.
- [18] **Vacaru, S. :** Nonholonomic Ricci flows and parametric deformations of the solitonic pp -waves and Schwarzschild solutions, *Electronic Journal of Theoretical Physics*, 6, N21 (2009), 63-93.